

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/78893/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Megardon, Geoffrey, Tandonnet, Christophe, Sumner, Petroc ORCID: <https://orcid.org/0000-0002-0536-0510> and Guillaume, Alain 2015. Limitations of short range Mexican hat connection for driving target selection in a 2D neural field: activity suppression and deviation from input stimuli. *Frontiers in Computational Neuroscience* 9 , 128. 10.3389/fncom.2015.00128 file

Publishers page: <http://dx.doi.org/10.3389/fncom.2015.00128>
<<http://dx.doi.org/10.3389/fncom.2015.00128>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Limitations of Short Range Mexican Hat Connection for Driving Target Selection in a 2D Neural Field: Activity Suppression and Deviation from Input Stimuli.

Geoffrey Mégardon^{1, 2*}, Christophe Tandonnet^{3, 4}, Petroc Sumner¹, Alain Guillaume^{2, 5}

¹School of Psychology, Cardiff University, United Kingdom, ²Aix-Marseille Université, France, ³Université de Genève, Switzerland, ⁴Aix-Marseille Université, France,

⁵Department of Psychology, New York University, USA

Submitted to Journal:

Frontiers in Computational Neuroscience

ISSN:

1662-5188

Article type:

Original Research Article

Received on:

26 Jun 2015

Accepted on:

02 Oct 2015

Provisional PDF published on:

02 Oct 2015

Frontiers website link:

www.frontiersin.org

Citation:

Mégardon G, Tandonnet C, Sumner P and Guillaume A (2015) Limitations of Short Range Mexican Hat Connection for Driving Target Selection in a 2D Neural Field: Activity Suppression and Deviation from Input Stimuli. *Front. Comput. Neurosci.* 9:128. doi:10.3389/fncom.2015.00128

Copyright statement:

© 2015 Mégardon, Tandonnet, Sumner and Guillaume. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](http://creativecommons.org/licenses/by/4.0/). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Full Title (155 characters/250):

Limitations of Short Range Mexican Hat Connection for Driving Target Selection in a 2D Neural Field: Activity Suppression and Deviation from Input Stimuli.

Short Title (50 characters/50):

Short Mexican Hat Limitations for Target Selection

Geoffrey Mégardon^{1,2,*}, Christophe Tandonnet^{3,4}, Petroc Sumner¹, Alain Guillaume^{2,5}

¹ : School of Psychology, Cardiff University, Tower Building, 70 Park Place, Cardiff, UK.

² : Laboratoire de Neurobiologie de la Cognition, UMR CNRS 6155, Aix-Marseille Université, 3 place Victor-Hugo 13331, Marseille, France.

³ : Faculté de Psychologie et des Sciences de l'Education, Université de Genève, 40 bd du Pont d'Arve, CH-1205 Genève, Suisse.

⁴ : Laboratoire de Psychologie Cognitive, UMR 7290, Centre National de la Recherche Scientifique, Aix-Marseille Université, 3 place Victor Hugo, 13331 Marseille, France

⁵ : Department of Psychology, New York University, 6 Washington Place, New York, NY, 10003, USA.

Corresponding author:

Geoffrey Mégardon, School of Psychology, Cardiff University, Tower Building, 70 Park Place, Cardiff, UK. E-mail: geoffrey.megardon@gmail.com.

Counts:

Abstract: 299 words

Main text (including references and figure legends): 10,494 words

Table: 1

Figures: 8

Keywords:

Dynamic neural field, action selection, center-surround, spiking neuron model, global effect, deviation away, saliency map

Abstract:

Dynamic Neural Field models (DNF) often use a kernel of connection with short range excitation and long range inhibition. This organization has been suggested as a model for brain structures or for artificial systems involved in winner-take-all processes such as saliency localisation, perceptual decision or target/action selection. A good example of such a DNF is the superior colliculus (SC), a key structure for eye movements. Recent results suggest that the superficial layers of the SC (SCs) exhibit relatively short range inhibition with a longer time constant than excitation. The aim of the present study was to further examine the properties of a DNF with such an inhibition pattern in the context of target selection. First we tested the effects of stimulus size and shape on when and where self-maintained clusters of firing neurons appeared, using three variants of the model. In each model variant, small stimuli led to rapid formation of a spiking cluster, a range of medium sizes led to the suppression of any activity on the network and hence to no target selection, while larger sizes led to delayed selection of multiple loci. Second, we tested the model with two stimuli separated by a varying distance. Again single, none, or multiple spiking clusters could occur, depending on distance and relative stimulus strength. For short distances, activity attracted towards the strongest stimulus, reminiscent of well-known behavioural data for saccadic eye movements, while for larger distances repulsion away from the second stimulus occurred. All these properties predicted by the model suggest that the SCs, or any other neural structure thought to implement a short range MH, is an imperfect winner-take-all system. Although those properties call for systematic testing, the discussion gathers neurophysiological and behavioural data suggesting that such properties are indeed present in target selection for saccadic eye movements.

1 Introduction

The ability to select important stimuli for further processing and action planning is a key function of brains of visually dominant animals. For instance, in primate visuo-motor systems only a small part of the retinal input benefits from a high spatial resolution; hence to select where to look is vital to extract relevant information from the environment. Points of interest have to be extracted from the overall visual input and, from those extracted points, only one can be selected at a time to orient gaze or attentional focus. Since Koch and Ullman (1985) it is thought that potential points of interest are evaluated through early visual processing and converge on a saliency map.

It has been suggested for a long time that a connectivity pattern of short range excitation and long range inhibition in topographically organized visual structures could achieve *saliency localization* - see blob detection models for computer vision (Bretzner and Lindeberg 1998; Kong, Akakin, and Sarma 2013; Lowe 1999) but also models of V1/LGN (Kang, Shelley, and Sompolinsky 2003; Schwabe et al. 2006; Spratling 2010; Zeng, Li, and Li 2011) - and *target selection* (Kuniharu Arai, Keller, and Edelman 1994; Kopecz 1995; Kopecz and Schöner 1995; Trappenberg et al. 2001). This connectivity pattern is often referred as a Mexican hat (MH) or center-surround inhibition, and was already implemented in early Dynamic Neural Field (DNF) models (e.g. Amari 1977). Recently the relevance of such organization has also been underlined for action selection in artificial cognition (Erlhagen and Bicho 2006; Richter, Sandamirskaya, and Schöner 2012; Sandamirskaya 2014); hardware implementations have emerged (Millner et al. 2010) and are suggested to be an important milestone for developing complex cognition (Indiveri, Chicca, and Douglas 2009).

Among neural structures often modelled using a DNF with MH connectivity (which we will refer to as DNF-MH), a prominent example is the superior colliculus (SC), a layered structure at the roof of the brainstem implicated in the control of gaze and attention orientation (Krauzlis, Liston, and Carello 2004; Munoz 2002; Robinson 1972; Sparks 1986; Sparks 2002; Guillaume and Péliesson 2001). The superficial layers of the SC (SCs) receive afferents directly from the retina and also from visual cortex and show strong visual activations. The intermediate-deep layers (SCi) display premotor activity for gaze orienting and receive multisensory input from a range of sources including connections from the SCs as well as 'top down' input from frontal cortex and basal ganglia. Both layers are topographically organized (retinotopic organization) and in register to one another. This neural structure is hence seen as a sensory-motor interface able to associate a motor command to visual information through connections between superficial and intermediate-deep layers (Isa 2002), as well as through other input. While both layers have been assumed to have MH connectivity, most modelling has focused on the SCi (and hence on target/action selection rather than saliency). Results of SCi studies (electrophysiology: McIlwain 1982, Munoz and Istvan 1998; and anatomy: Behan and Kime 1996; Meredith and Ramoa 1998) were in favour of MH connectivity and also suggested that inhibition from a given site can concern very distant areas of the map. Hence, without more precise measures, it was assumed that the inhibitory influence was very large. Numerous models implementing long range inhibition (K. Arai, Keller, and Edelman 1993; Bompas and Sumner 2011; Kopecz 1995; Kopecz and Schöner 1995; Marino et al. 2012; Meeter, Stigchel, and Theeuwes 2010; Trappenberg et al. 2001; Wilimzig, Schneider, and Schöner 2006) showed that it was successful for winner-take-all selection of a saccade target among several options.

However, the idea of long range inhibition in the SC has been challenged (Isa and Hall 2009; Lee and Hall 2006). Very recently, a clearer picture has been obtained. Phongphanphanee et al. (2014) using multi-electrode arrays on slice preparations of rodent SC evaluated the local connectivity in SCs and SCi. This study found MH connectivity only in SCs and that, in this case, the range of inhibition is relatively short (see below for details). In SCi, the excitation zone was at least as large as the area of inhibitory influence. The second main difference between the SCs and the SCi revealed by the study concerned the time course of their excitatory and inhibitory responses to a sustained stimulation:

where the SCi was behaving as an accumulator, the SCs showed transient responses. Globally these results led the authors to conclude that the winner-take-all phenomenon is observed in the SCs and that it enables saliency detection. Phongphanphanee et al. (2014) wrote "The sensory layer (SCs) is optimized to localize the single most salient stimulus" (p. 2342). The SCi, in turn, would cascade activity from the SCs and integrate it with its other inputs to perform target selection. Importantly, the saliency selected and localised by the SCs can be translated into the winner of the SCi target selection when other target candidates are negligible. As stated above, numerous models of the SC were implementing long range inhibition to perform selection. The results of Phongphanphanee et al. (2014) call for an exploration of properties of map integrating MH with short range inhibition and temporal dynamics based on the SCs.

The aims of the present study were: 1) to test the capacity of such a DNF-MH with short range inhibition to perform reliable target selection and 2) to highlight its noticeable properties and its potential limitations in such a context. Importantly, those properties could represent testable predictions to address if the SCs – or any brain structure – performances are indeed driven by a short range MH. We implemented this type of DNF-MH in two dimensions with spiking neurons. We fed it with various types of input stimulation to assess the emergence of localized and stable clusters of firing neurons (a "spiking cluster") that would represent saliency and/or target selection. We first explore the effect of stimulus size on the performance of the model. Second, we tested the model while two stimuli were presented at the same time and we measured their interaction while varying their weights and the distance between them.

To anticipate some of the key results, varying the size of a single stimulation led to bimodal activation and to centre-surround interactions that could result in the complete suppression of any activity on the network. When two stimulations were used, phenomena of attraction, complete suppression, and repulsion were observed for different distances. Applied to target selection, those properties may represent detrimental phenomena: prior loci of interest extracted from feature maps could suppress themselves, or produce clusters of activity that are not localised on the stimuli of interest. Interestingly, we can link these properties with previous neurophysiological and behavioural studies. These links are extensively explored in discussion.

As a final note, although our rational is largely based on results obtained in the SC, especially the superficial layers, our results describe a set of phenomena possible for any DNF-MH implementation and usage.

2 Material and Methods

2.1 Overview of the Model

The model is a simple network of neurons organized as one 2D layer of 100x100 cells (Figure 1A) and connected according to a 2D Mexican hat kernel (Figure 1B). Our model is close to those of K. Arai, Keller, and Edelman 1993; Marino et al. 2012; Trappenberg et al. 2001; Wilimzig, Schneider, and Schöner 2006. Nevertheless the critical differences are that we implemented a MH with a short range of inhibition and spiking neurons (Figure 1C) which allow to set up different synaptic decay times for inhibition and excitation (see section 2.2). Finally, we did not implement the logarithmic compression of space that is observed in the SC to remain general.

The model is implemented in Python 2.7 (<http://www.python.org/>) and using the library BRIAN, a spiking neuron network simulator (D. Goodman and Brette 2008; D. F. Goodman and Brette 2009). The code source for all the following simulations can be found at: <https://github.com/Nodragem/SuppData-MHLimitations-Selection>.




Figure 1: Overview of the model. **Subplot A:** The model is a dynamic neural field (DNF) of 100x100 cells. The red to yellow circle represents the cluster of spiking neurons after stimulation of the neurons on the green line fed by the input neuron (green circle). This cluster forms a circle centred on neuron A. Neurons A (blue-white dot) and B (red-white dot) are marked in reference to subplots B and C. **Subplot B:** Illustration of the Mexican hat kernel. The graph shows the connection weight of neuron A with its neighbourhood. The X- and Y-axis represent the distance from neuron A in number of cells; The Z-axis represents the weight of connection, a positive number is excitatory while a negative number is inhibitory (arbitrary unit). **Subplot C:** illustration of equation 1. Each spike of neuron A (panel 1, red bars) opens excitatory channels on the membrane of neuron B that close by themselves according to time constant τ_e (panel 2, orange curve). These opened excitatory channels raise the membrane potential of neuron B (panel 3, green curve). When a threshold (-50 mV here) is reached, a spike is triggered in the neuron B (panel 4, red bar).

The spiking neuron model (Lapicque 1907; Brunel and van Rossum 2007) used here is a simplification of conductance-based integrate-and-fire (Hodgkin and Huxley 1952; Shadlen and Newsome 1998). Activity of each neuron (unit) of this network can be described with the following equations:

$$\left\{ \begin{array}{l}
\tau_m \frac{\partial V(n, t)}{\partial t} = -(V - V_0) - g_e(V - V_e) - g_i(V - V_i) \quad (a) \\
\frac{\partial g_e(n, t)}{\partial t} = -\frac{g_e}{\tau_e} + \alpha_e \sum_{n'} \sum_f \delta(t_{n'}^f - t) w_e(n', n) + \alpha_s w_s F_s \quad (b) \\
\frac{\partial g_i(n, t)}{\partial t} = -\frac{g_i}{\tau_i} + \alpha_i \sum_{n'} \sum_f \delta(t_{n'}^f - t) w_i(n', n) \quad (c)
\end{array} \right. \quad (1)$$

with $V = V_r$ if $V > V_t$

168

169 **Equation (a)** describes the time course of the membrane potential V for all neuron n (Figure 1C,
170 column 3). It goes toward its equilibrium V_0 with a time constant τ_m when at rest while it goes toward
171 V_e or V_i when g_e or g_i are different from zero. When V reaches a threshold V_t for a neuron n , a spike
172 is emitted and V is reset to V_r for that neuron n . After it emits a spike, a neuron will be unaffected by
173 any input during a refractory period of 1.5 ms. This refractory period limits the maximum firing rate
174 to 600 Hz which is consistent with SC cell recordings for instance (Anderson et al. 1998; Sparks,
175 Holland, and Guthrie 1976).

176 **Equations (b) and (c)** describe the time course of the opening of excitatory and inhibitory gates --
177 g_e and g_i -- on neuron n 's membrane (Figure 1C, column 2). By default, g_e (respectively g_i) goes to
178 zero with a time constant of τ_e (respectively τ_i) -- the synapse decay time. For each time $t_{n'}^f$ --
179 corresponding to a spike f of a neuron n' in the network -- g_e (respectively g_i) gets an immediate
180 increase which corresponds to the weight connecting n to n' defined by w_e (respectively w_i). Finally,
181 one or more experimenter-controlled spiking neurons can be connected to the model through g_e (see
182 Figure 1A). Their firing rate over time is controlled by a curve F_s ; in that sense, they resemble electric
183 stimulations used in neurophysiology and do not follow a Poisson process. The connections to the
184 network are defined within the unit interval with a matrix w_s and are modulated with α_s .

185 The matrices of connections w_e and w_i are normalized between -1 up to 1: α_e and α_i are used to scale
186 them to a relevant dimension for the network, its unit being millivolt. The matrices w_i and w_e are
187 computed from a difference of Gaussians equation:

$$f(x, y | \theta_x, \theta_y, K, \beta) =$$

$$(1 + \beta) \exp \left(-\frac{(x - \mu_x)^2}{2\sigma_x^2} - \frac{(y - \mu_y)^2}{2\sigma_y^2} \right) - \beta \exp \left(-\frac{(x - \mu_x)^2}{2K^2\sigma_x^2} - \frac{(y - \mu_y)^2}{2K^2\sigma_y^2} \right)$$

(2)

190 The resulting subtraction gives a Mexican hat curve (see Figure 1B and Figure 2); the first term on
191 the right hand side of equation (2) is used as w_e and the second term as w_i . The variables σ_x/σ_y and
192 $K\sigma_x/K\sigma_y$ define the standard deviation of the Gaussians. Thus, K is used to set up the
193 inhibitory:excitatory extent ratio. β is a parameter controlling the depth of the inhibition.

194

2.2 Parameter Choice

All the parameters of the model are summarized in the Table 1. Parameters for which values are not given in the table have values varying in the different simulations and these values are to be found in the description of each specific simulation. The neuron parameters were taken from recent spiking neuron models of the SC (Lo et al. 2009; Morén, Shibata, and Doya 2010; Morén, Shibata, and Doya 2013) and adapted to obtain clusters that maintain a stable activity on the map for the range of stimulations we used. Concerning synaptic decay times, τ_i is superior to τ_e which is coherent with the observation of Phongphanphane et al. (2014) during a sustained stimulation of the SC (see their figure 7): indeed, Figure 5E of the present report shows that our model was able to reproduce a transient response of the membrane potential to a sustained input. However, to our knowledge the time constants τ_i and τ_e of actual SC neurons have never been specifically measured, which explains large differences in parameters values between the work of the two previous teams. By default, no noise will be introduced in the model. If a noise source is used, it will be stated in the text.

Concerning the lateral connection parameters, we used values for K and β that were chosen based on previous physiological or modelling studies. K , corresponding to the ratio inhibition-extent/excitation-extent, was set to 1.2 to limit lateral inhibitory influence to a relatively small range consistent with recent results (see Isa and Hall 2009 for a review). This ratio is similar to the value suggested by the SCs in-vitro study of Phongphanphane et al. (2014). Indeed, they reported an EPSC half-width area of $130 \mu\text{m}^2$ and IPSC half-width area of $145 \mu\text{m}^2$ (see their figure 4.D and their text page 5; note also that Lee and Hall's (2006) *in vitro* study on rat SCi reported ratios of $500\mu\text{m}/300\mu\text{m} = 1.6$ or $500\mu\text{m}/400\mu\text{m} = 1.25$) β corresponds to the strength of inhibition; it was set at 6.0 in order to set the maximum inhibition weight at roughly the half of the excitation maximum weight to fit with the results of Arai et al. (1994) (see the black curve of our Figure 2 -- the minimum weight of the reference MH is at -100 mV for a maximum of 200 mV). Note that we test variations in these K and β values below.

Lastly, our parameters are chosen for the neural field to be bistable between the all-off state and a spiking cluster state. When a bump in the membrane potential reaches the threshold, the model generates systematically a stable and well-defined group of spiking neurons around the point which passed the threshold. We name this group a "spiking cluster" to distinguish it from bumps in the membrane potential. This spiking cluster is similar to a bump of activity in a population rate model, and being stable, it survives after we stop stimulating the neural field.

Table 1: Model parameters and variables:

| General parameters | Symbol | Value and unit |
|---------------------------------------|------------|-----------------|
| simulation time | None | 200 ms |
| map size | None | 100x100 neurons |
| simulation clock precision | None | 0.01 ms |
| recording clock precision | None | 1 ms |
| Neuron parameters | Symbol | Value and unit |
| membrane time constant | τ_m | 10 ms |
| excitation time constant | τ_e | 3 ms |
| inhibition time constant | τ_i | 10 ms |
| potential threshold | V_t | -50 mV |
| reset potential | V_r | -80 mV |
| resting potential | V_0 | -70 mV |
| Nernst potential of excitation ions | V_e | 0 mV |
| Nernst potential of inhibition ions | V_i | -80 mV |
| Neuron variables | Symbol | Unit |
| membrane potential | V | mV |
| number of opened excitatory channels | g_e | no unit |
| number of opened inhibitory channels | g_i | no unit |
| Mexican hat parameters | Symbol | Unit |
| depth of inhibition controller | β | no unit |
| Inhibition/excitation extent ratio | K | no unit |
| standard deviation on Y-axis | σ_y | cells |
| standard deviation on X-axis | σ_x | cells |
| center position on X-axis | μ_x | cells |
| center position on Y-axis | μ_y | cells |
| matrix of positive connections | w_e | no unit |
| weight factor for positive connection | α_e | 200 mV |
| matrix of negative connections | w_i | no unit |
| weight factor for negative connection | α_i | 200 mV |
| External stimulus parameters | Symbol | Unit |
| spikes train | F_s | no unit |
| matrix of connections with the model | w_s | no unit |
| weight factor | α_s | mV |

239

240

241

242

243

244

245

2.3 Simulation Set 1: Size Variation of a Single Stimulus

In a first set of simulations we want to characterize the response of the selection map to stimuli of different sizes. We iterate the exploration for three different instances of the MH to test for the effect of slight variations in the inhibition strength and extent.

A range of stimulus lines of varying size (see below) was tested with the model. We ran three different sub-simulations (S1, S2, S3): one testing the reference MH (see general method) and two testing variations of it in order to make sure that the results are robust to moderate changes in the connectivity profile. The first sub-simulation (S1) implemented the reference MH ($K = 1.2$ and $\beta = 6.0$; see Model parameters). The second sub-simulation (S2) was conducted to test a larger extent of inhibition. K was fixed to 2.0, which covers the upper end suggested by data of Phongphanphanee et al. (2014). In order to only address the extent of inhibition, β was set to 1.43 with an optimization algorithm to keep the minimum of MH function (depth of the inhibition) similar to S1. The third sub-simulation (S3) was conducted with $K = 1.2$ and $\beta = 8.0$ to observe the effect of a stronger inhibition while keeping its extent constant. For the whole set of simulations in this part, σ_x and σ_y were fixed to 5 neurons, this was chosen to get relatively small MH lateral connections compared to the dimension of the model map. Given that the determinant factor is the relative size of the stimuli compared to the MH's size, having small connections allowed us to increase the range of tested stimulus sizes.

The map was stimulated with line-shaped stimuli with twenty sizes (2 neurons up to 42 neurons in steps of 2 neurons along the Y-axis). These line-shaped stimuli were defined by $I_s = \alpha_s * w_s * F_s$ as explained in the equation (1.b). The maximum size of 42 neurons represents less than 50% of the Y-axis size of the model map in order to limit border effects (map size = 100x100 neurons, see table 1). The firing rate pattern over time of the external input, F_s , was a Gaussian centred on 25 ms with a standard deviation of 80 ms and a maximum frequency of 400 Hz. The strength of the stimulus was $\alpha_s = 4000\text{mV}$. Finally, the duration of each simulation was 200 ms.




Figure 2: The three Mexican hats tested in Simulation 1. They correspond to the connection of a neuron n with its neighbourhood (i.e. $ae.we + ai.wi$). They are only plotted on the X-axis and for one direction.

The results will be split in four parts. The first part focuses on the spatial pattern obtained on the map, the second part on the temporal dynamic. The third part investigates if our result would be different if using a sustained input ($F_s = 400\text{Hz}$) instead of the Gaussian firing rate pattern aforementioned. Finally the last part extends our results to bidimensional shape – replacing the line-shaped stimuli by squares and circles.

2.4 Simulation Set 2: Interaction of Two Stimuli

Our first set of simulations addresses the effect of stimulus size in a simple DNF-MH used as a target selection map. However, such a map is prone to receive many candidate points of interest from satellite structures feeding it. Our second set of simulations tests the behaviour of our DNF-MH model when stimulated at two points with varying the distance and relative strength. In a comparison of our model with the SCs, this simulation is analogous to the in-vitro experiments conducted by Phongphanphanee et al. (2014) and by Vokoun et al. (2014) in which these two teams stimulated two points in the SCs varying the distance and the strength of stimulations injected in each point.

Two stimulation points, A and B, of size 2×2 , are considered. Stimulation A is kept at a fixed location ($x = 31$; $y = 51$), while stimulation B is tested for distances from 2 to 40 cells with a step of 2. Stimulation A and B both have the same firing rate pattern as used in simulation 1. While the stimulation B is always connected with a weight of 4000 mV to the model, stimulation A is tested for 3 different weights: 1333 mV, 2000 mV and 4000 mV. We used the reference MH configuration ($K = 1.2$; $\beta = 6.0$) but in a larger implementation ($\sigma_x = \sigma_y = 8.5$ cells, compared to 5 cells in Simulation 1, to increase the MH size and hence virtually increase the granularity of our probing). The result we report here is the position of the spiking cluster nearest to stimulation B on the map. Its localization is defined by the centre of gravity of its spike count over all the simulation. To control that border effects was not at the origin of the following observations, a control condition was run that tested the spiking cluster position for the different location of the stimulation B alone. The spiking cluster positions were then well aligned with the stimulation B and suggest there is not border effect at those locations.

The results for the simulation set 2 will be split in two parts. The first part presents the results without including noise in the model. The second part tests if the results obtained for the condition 4000mV-4000mV are robust to the addition of noise in the model and if they extend to a slight inequality in A and B intensity (3500mV-4000mV). Precisely, the noise was added to the membrane potential and was following a normal distribution of standard deviation 4 mV.

3 Results

3.1 Simulation Set 1: Spatial Patterns

Figure 3 shows membrane potential and firing rate for all neurons of the neural field for a subset of stimulus sizes for the 3 MH variants (S1, S2, S3, depicted in Figure 2). These values of membrane potential and firing rate were averaged over all the simulation time (200ms). The represented line sizes illustrate the different observed activity patterns. Panel D shows the number of spiking clusters (see parameter section for definition) as a function of the stimulus size for the three MH variants, as further explained below.




Figure 3: Overview of the spiking clusters spatial distribution during S1, S2 and S3 (respectively subplot A, B and C). The results are shown for the most informative stimulus sizes, which are different according to the set of simulations and are indicated on each column of the graphs. **Subplot A, B, C:** On each picture, the top part shows average membrane potential during the simulation; the lower part shows average firing rate during the simulation. **Subplot D** summarizes the result: the number of spike clusters is computed as the sum of spikes on the map divided by the sum of spikes occurring for the first distractor size (each first distractor size gave rise to one spike cluster that is used as reference).

For the first MH (S1, $K = 1.2$ and $\beta = 6.0$, Figure 3A and black line in 3D), a unique circular spiking cluster located on the centre of the stimulus line was observed from the smallest size up to the size of 18. Despite the transient nature of the stimulation, the spike cluster persists during the whole duration of the simulation. On the contrary, from size 20 to the largest tested size (size 42), no spiking clusters appeared on the firing rate map: a complete activity suppression was observed. It can be noticed that on the membrane potential map, the activity appears equally spread for size 20 while the activity is stronger on the extremities for size 42. This sub-threshold activity distribution suggests that the extremities could win the competition if the threshold was decreased.

In the second sub-simulation (S2, $K = 2.0$ and $\beta = 1.43$, Figure 3B and red line in D), we observed similar results but the spiking cluster for small line sizes was larger and the complete activity suppression starts at a larger stimulus size. These two observations are to be related to the slightly larger excitation influence in S2 with respect to S1 (Figure 3). The main difference with S1 appears at size 42: the activity on the extremities was strong enough to give rise to two spiking clusters. Those two spiking clusters have a weaker average firing rate than the one observed for previous sizes; below we show this is due to a larger latency before the first spike rather than a lower firing rate once initiated (see next section).


In the third sub-simulation (S3, $K = 1.2$ and $\beta = 8.0$, Figure 3C and green line in D), similar results as in S2 and S1 are found but with a smaller radius of the spiking cluster for small line sizes and a suppression that starts at a smaller stimulus size (size 16). Here, two spiking clusters were observed, as in S2, for stimulus size 30 to 36. However, S3 differs from S2 as a complete suppression was again observed for larger sizes than 36. When present, the two spiking clusters were on average weaker than the unique spiking cluster observed for smaller sizes, and again this is due to delay rather than firing rate once the spiking cluster occurs (see below).

Thus complete activity suppression occurred for at least one range of sizes for each set of simulations. The stimulus size for which it appears is positively correlated with the size of the positive area of the MH used for these simulations.

Lastly, note that the stimuli tested were spatially homogenous: each point of the stimulus gave the same input to the map. This type of stimulation may favour complete suppression, and if noise were present in the network, it is conceivable that it could randomly favour the selection of a spiking cluster and hence eliminate the phenomenon of complete suppression. To test this hypothesis we added normally distributed noise in the membrane potential of all the units of the 2D network, using $K = 1.2$ and $\beta = 6.0$ (S1). The standard deviation of the noise was of 4mV, which corresponds to a fifth of the distance between the resting potential and the threshold. The results were similar to those presented above. Hence, even with noise in the network, the phenomenon of complete suppression could be observed.

3.2 Simulation Set 1: Temporal Dynamics

In simulations S2 and S3 larger stimuli could lead to two spiking clusters, which show a lower firing rate average than for the unique cluster appearing for smaller sizes. Figure 4A, B, C shows the evolution of the membrane potential for neurons just next to the stimulus line (see caption for more details) for the same sizes addressed in Figure 3A, B and C. It can be observed that the threshold to the first spike is reached much later for sizes giving rise to two spiking clusters (size 42 in SA2 and size 30 in SA3) when compared to sizes leading to one spiking cluster. In addition we have estimated the firing rate of the spiking clusters for the last 50 ms of each simulation: their firing rate does not change between stimulus size (550-600Hz for S2, 350-400Hz for S3,). Hence the change in firing rate average observed in S2 and S3 was the result of the change in latency for the membrane potential to reach the threshold.



386

387 Figure 4: Overview of the membrane potential dynamics during SA1, SA2 and SA3. **Subplot A-C:** Effect of the stimulation on the
 388 neighbourhood according to time. We report the membrane potential of the most excited neuron among the neurons situated along a
 389 line parallel to the stimuli and at 2 cells from it ($x = 52$). When a neuron on this line reaches its threshold (at -50 mV, see the dashed
 390 horizontal line), it means that a spiking cluster is created. **A, B and C** correspond to the 3 different MHs introduced in figure 3; their
 391 parameters K and β (from equation 2) are indicated. For each MH, we plot the maximum of the membrane potential for the sizes shown
 392 in figure 4. The vertical lines at the bottom represent the input spike train. **Subplot D:** Initial speed (averaged between 0 and 6 ms) of
 393 the membrane potential (in mV/s) according to stimulus size for SA1, SA2 and SA3. The circles plotted on the curves denote that, for
 394 these stimulus sizes, the membrane potential reached the threshold before 30 ms and led to one spiking cluster on the neural field.
 395 **Subplot E:** Speed of the membrane potential (in mV/s) between 30 and 90 ms when auto-inhibition prevented threshold being reached
 396 in the first rise, plotted according to stimulus size. The circles plotted on the curves denote that, for these sizes, the membrane potential
 397 reached the threshold sometime after 30 ms and there are two spiking clusters on the map

398 We can observe in all the simulations (Figure 4A, B, C, all curves) an early rise of membrane
 399 potential. This early rise is at the origin of all single spiking clusters observed in Figure 3. Figure 4D
 400 shows the speed of this early rise (for the interval between 0 and 6 ms) for all stimulus sizes. This
 401 speed increases until an optimal size (10, 12 and 8 cells for S1, S2 and S3 respectively) and then
 402 decreases to a plateau. The obtained curve is analogue to what is found with the firing rate of neurons
 403 in surround suppression literature (Sceniak et al. 1999; Schwabe et al. 2006). Empty circles on the
 404 curves indicate that the threshold is reached before 30 ms (i.e. a single spiking cluster is observed).
 405 Hence we can see that close to the stimulus size corresponding to the beginning of the plateau, the
 406 initial wave of excitation starts to fail to reach spiking threshold. Interestingly, in those conditions
 407 (size 20 and 42 of Figure 4A for instance), we can observe that the early rise is transient. This transient
 408 nature will be explained below with Figure 5 showing the dynamics of excitatory and inhibitory
 409 influences. Interestingly, this transient rise in the membrane potential echoes the transience observed
 410 by Phongphanphanee et al. (2014) in the SCs as previously mentioned (see their figure 7A, left).

411 The two spiking clusters for larger stimuli occurred through a late second rise in membrane potential
 412 after 50ms (e.g. size 42 and 30 for S2 and S3). By observing the curve for size 20 in S1 Figure 4A,
 413 we can see that a late rise in the membrane potential occurs also for intermediate sizes, but
 414 insufficiently to produce late spiking clusters (see also size 24 for S2 and sizes 16 and 38 for SA3),
 415 this corresponds to the complete activity suppression shown on Figure 3. To examine this further,
 416 Figure 4E plots the mean speed of the membrane potential averaged between 30ms and 90ms to
 417 illustrate how the second rise varies over stimulus size. The time window used catches the variation

for S3 especially well showing that the second rise of the membrane potential, like the first, also has an optimal stimulus size after which the rise speed decreases again and it fails to reach threshold (compare with the panel D of the Fig 3).

3.3 Simulation Set 1: Effect of Input Dynamic

To get a better view of the dynamics of inhibition and excitation, we compared these dynamics for a sustained input at 400Hz to those for the transient input with a Gaussian profile as previously used. Figure 5 shows the membrane potential of the neuron showing the largest hyperpolarization near the stimulus in these two cases (panels A,B,C for transient input and panels D,E,F for sustained input) obtained with parameters of S3 ($K = 1.2$, $\beta = 8.0$). It highlights that the dynamic of the initial transient rise in membrane potential comes from a delayed wave of inhibition (Panels B, C, E and F). Indeed, the weight of excitation is twice larger than the weight of inhibition, giving an initial advantage to the excitation which the inhibition later catches up due to its larger decay time constant. Note that this wave of inhibition comes from remote units (see Figure 2) and, so, does not appear for small size (Panels A and D).

With the sustained input, we still observe a second rise in the membrane potential; the inhibition curve still decreases after having overtaken the excitation. As our recording takes place toward the extremities of the stimulus -- near the potential winner loci -- this decrease of inhibition thus comes from the decrease of activity of the middle of the stimulus (see Figure 3B size 42, and 4C size 30, 38) silently losing the competition at sub-threshold level.

For a stimulus size of 36, no spike cluster is observed in this sustained input condition in opposition with the transient stimulus condition (Figure 5.C and 5.F). As mentioned above, the second membrane potential rise is weaker when using the sustained stimulation. Counter-intuitively, this suggests that to decrease or to stop the stimulation input --with the transient stimulation-- helped the membrane potential to reach the threshold. Two pieces of explanation are that 1) as the neurons are excitatory coupled, the most excited regions of the stimuli self-sustain their firing longer than the others when our input stops, 2) the most inhibited regions lose their only source of excitation when our input stops. Then to stop or decrease the input signal can accentuate disequilibrium in the competition and facilitate a target selection outcome.




Figure 5: Excitatory and Inhibitory Channels opening and membrane potential according to time for transient stimulation (upper graphs) and sustained stimulation (lower graphs). Data from SA3 ($K=1.2$, $\beta=8.0$) are shown here; similar curves can be obtained from the other conditions. The transient input (A,B,C) is the one used in the previous simulations and corresponds to a Gaussian (see F_s in Inputs and Methodology). The sustained input (D,E,F) set $F_s = 400$ Hz, e.g. the firing rate of the input stimulation is at 400 Hz and constant over the time. The vertical lines at the bottom represent the input spike train. The tracked neurons all come from the row at 2 cells from the stimulated neurons ($x=52$). At each timepoint, the following measures are extracted from the neuron that has the maximum membrane potential among the tracked neurons. The curve “V” is the evolution of the membrane potential over the time. “ge” or “gi” describe the evolution of, respectively, the number of excitatory or inhibitory opened channels on the neuron’s membrane. However, for the sake of comparison, the number of opened channels ge and gi are multiplied by a scaling factor. Indeed, for any value of V: $|E_e - V| > |E_i - V|$ where E_i and E_e are, respectively, the inhibition and the excitation equilibrium. Thus, an excitatory gate that opens always has more effect on V than an inhibitory gate. The scaling factors represent this difference by being $|E_e - \hat{V}|$ for ge and $|E_i - \hat{V}|$ for gi, with $\hat{V} = (V_t - V_0)/2$.

3.4 Simulation Set 1: Generalization to 2D stimulus shapes

Our DNF-MH model of target selection map shows a phenomenon of total activity suppression related to stimulus size for 1D stimulus. Here, we generalize our observations to 2D stimuli by testing the behaviour of the model when stimulated with a circle and a rectangle of varying size. We used the reference MH ($S1$, $K = 1.2$ and $\beta = 6.0$).

Figure 6 shows results obtained for these tests conducted with 2D shapes. Columns 1 and 3 show average firing rate over the simulation period and columns 2 and 4 contain the spikes train of neurons on the diagonal of the map. Similar phenomena of activity suppression to those for the 1D stimulus are observed: spiking clusters did not emerge for size 18 for the square and for sizes 20-26 for the circle. Additionally for further increases of size, several clusters appear: from sizes 20 (square) and 28 (circle) four spiking clusters emerged (Note that activity for the circle segregates into 4 regions because the pixelation of our map, theoretically no point on a disk would be advantaged on a continuous competition field). Especially in the case of the circle, these clusters tend to move as if they are repulsed from the centre (lower panel of the column 4). This repulsion becomes weaker with size until a new spiking cluster emerges at the centre in addition of the four on its corners (not shown).

The spike trains (columns 2 and 4) also allow observing a latency increase for clusters appearance when increasing the size. For smaller sizes, below 18 (square) and 20 (circle), the unique cluster appears with almost no delay with respect to the onset of the stimulation. Conversely, when spikes appear for larger sizes, whether they finally disappear (size of 18 for the square or of 20-26 for the circle) or are part of stable spiking clusters (larger sizes), there is a short latency period of

approximately 10 ms before their appearance. This latency increase is similar to the one observed when a 1D stimulus resulted in two clusters (see above) but of lower value: the latency increase for the 1D stimulus was around 70 ms. Finally, the first burst of spikes and the following gap (just at the beginning of the stimulation; middle and lower panels in the columns 2 and 4) can be respectively related to the initial rise of membrane potential and to the wave of inhibition seen with 1D stimuli. These differences can be explained by the greater number of neurons interacting, which speeds and strengthens excitatory and inhibitory influence. Hence, apart from this difference in the latency, results for 2D shapes are similar to those obtained for the 1D stimulus (line).




Figure 6: Results of the simulations for the Simulation 2 which test for a suppression effect with a square and a circular shaped stimuli. Column 1 and 3 show the average spike frequency (firing rate) in the neural map in hertz. The column 2 and 4 show the spikes train of neurons according to time. The red vertical lines at the bottom represent the input spike train. The recorded neurons are those forming the diagonal of the neural field from the position (25, 25) to the position (75, 75).

3.5 Simulation Set 2: Spatial Interactions

Panel B of Fig 7 shows a summary of the results for these simulations. The spiking clusters produced by the model indicate which locus has been selected as a target. Its deviation from stimulation B is shown as a function of distance between the two loci of stimulation. Negative values correspond to a deviation toward the locus of A. The three curves correspond to the three different intensity of stimulation tested for the point A (1333, 2000, 4000mV; B is always stimulated with 4000mV). Filled symbols indicate that only one spiking cluster was present on the network and open symbols that two clusters survived.

508 In the case of equal strength for both stimulations (green curve), for the first distances up to 14 cells,
509 we observed one single resulting cluster (fusion phenomenon) that was in between the two stimulation
510 loci (see panel A1 Attraction). That observation can be related to the activation merging found by
511 Vokoun et al. (2014) in the SCs (see their figure 3). Then for two following distances (16 and 18
512 cells), a complete suppression of activity on the map was observed (see panel A2 Suppression).
513 Finally, from a distance of 20 cells up to the largest tested distance (40 cells), two clusters are
514 produced and the closer to the site of B is repulsed in the opposite direction with respect to A (positive
515 values on y axis of the panel B). The panel A3 allows us to see that the same repulsion was observed
516 for the cluster close to the A site. This repulsion phenomenon decreased as the distance between the
517 two stimulating sites increased.

518 When stimulation B was stronger than stimulation A (blue and red curves), in almost every case only
519 one cluster was produced: a winner-take-all mechanism occurred and selected a locus near stimulation
520 B. Nevertheless, a deviation toward stimulation A is still observed up to the distance 16 cells: the
521 spiking cluster appears in between the two stimulations. Note here that the selected locus is closer to
522 the strongest stimulation and that it gets closer when the latter gets stronger. That bias toward the
523 strongest stimulus is also observed in Vokoun et al. (2014). For larger separation distances, the
524 winning cluster remained localized near site B. This result goes in line with the results of
525 Phongphanphanee et al. (2014): when the stimulations are close enough, an activation is present at A
526 and B sites while when the stimulations are more distant, no activity is recorded close to the
527 stimulation A (the weakest) and a normal cluster is observed close to the stimulation B (the strongest).
528 Nevertheless, the winner-take-all mechanism is not perfect: the selected locus is near to B but not
529 aligned with it. Indeed a deviation away from stimulation A occurred, similar to what we observed
530 with equal strength simulations.

531 Panels C1 and C2 show this winner-take-all phenomenon. However, for the last tested distance with
532 $A=2000\text{mV}$ (40 cells), the activity at A escaped from the inhibition influence of the stimulation B
533 and two clusters emerged (panel C3) which may be seen as a fail to select one target from the two
534 input: the stimulation A overcomes the surround inhibition – which decrease with the distance in that
535 range of distances – and stimulation B gives rise to its own spiking cluster. That does not occur for
536 the condition $4000\text{mV}-1333\text{mV}$, where the stimulation A is too weak to overcome the inhibition even
537 at such a distance.





Figure 7: Interaction between two stimulation induced bumps according to their distances. The magenta dot in subplot A and C represent the position of stimulation B, while the green dot represents the stimulation A. The white dot in subplot A is the centre of gravity of the spiking cluster the nearest from stimulation B. The plot B describes the deviation of that centre of gravity (white dot) from the stimulation B (magenta dot) on the x-axis. Filled dots denote there is only one spiking cluster on the map, while the unfilled dots denote there are two spiking clusters on the map. The simulation was run for different distance between the stimulation A and B (x-axis), and for different strength of the stimulation A (curves red, blue and green). Note that the subplot A shows an average of the firing rate over the simulation while the subplot B show an average of the membrane potential over the simulation.

3.6 Simulation Set 2: Tight competition and addition of noise.

Our previous results for two stimulation inputs of exactly same strength show that for a given range of distances a complete suppression of activity is observed. This corresponds to a failure for the DNF-MH model to select only one target. One can suggest that this failure of the winner-take-all is due to 1) the absence of noise in our model or 2) the unnatural exact equality of the two stimulations in competition.

554 We tested here if the previous results, notably the suppression, can be obtained with the addition of
555 noise and for close competition (3500mV vs 4000mV).

556 The Figure 8 shows the results of one simulation with these conditions. The results are strongly
557 similar to those obtained in the simulation without noise. The addition of noise, even if it helps to get
558 only one winner (compare distance 20 and 22 in Figure 8 and Figure 7B), does not prevent the case
559 of two-winner and no-winners in the 4000mV-4000mV condition. Interestingly, the condition 4000-
560 3500mV (dark blue curve) shows that we can also obtain activity suppression (Figure 8) when the
561 two stimulations are not exactly of the same strength. This occurs for distance 16 and 18 cells,
562 similarly to the equal strengths condition. However, in this case, the two-winner situation is not
563 observed directly after the suppression phase. For some distances, the curve is similar to the one
564 obtained in the condition 4000-2000mV. Nevertheless, finally stimulation A succeeds to give rise to
565 a spiking cluster because the inhibition from stimulation B gets smaller after a certain distance (refer
566 to the shape of a MH curve, Figure 2). Here, stimulation A being stronger than in the 4000-2000mV
567 condition, it overcomes the inhibition of B (i.e. it results in two clusters) at a smaller distance.



568
569 *Figure 8: Interaction between two stimulations according to their distances, with noise, and with an additional condition (4000-*
570 *3500mV) testing for tight competition. Same description as for Figure 7B.*

571

572

4 Discussion

The aim of the present study was to get further insight into the properties -- and their consequences for saliency or target selection -- of Dynamic Neural Fields based on a Mexican hat kernel (DNH-MHs) in the specific case of short inhibitory influence. Indeed, this type of lateral connection has been recently demonstrated for a classical biological example of DNF-MHs, the superior colliculus (SC; see Introduction). We designed a simple one layer model implementing the most recent data concerning lateral interaction in this neural structure (Phongphanphanee et al., 2014) and we tested its properties. We observed that certain stimulus sizes could lead through centre-surround interactions to a complete suppression of the network activity, while larger sizes led to multi loci selection. This complete suppression, which led to no target selection, also occurred when two stimulations were presented simultaneously within a certain range of distances. For smaller distances, the model selected a position in-between, closer to the strongest stimulus (attraction/fusion), while for larger distances the model selected two loci that were deviated away from the stimuli positions (repulsion). We discuss these results of suppression and spatial deviation (i.e., attraction/fusion and repulsion) obtained here in view of neurophysiological, modelling and behavioural previous findings.

4.1 Suppression phenomena: Neurophysiology results

It may seem counterintuitive to observe complete suppression on a saliency map for large stimuli of interest. This result can nevertheless be related to previous neurophysiological, modelling and behavioural findings in the visuo-oculomotor system and may help to disentangle unanswered questions.

Suppression phenomena in which larger stimuli produce lower activity than smaller ones are well described in sensory systems and especially in the visual system (Allman, Miezin, and McGuinness 1985; Series, Lorenceau, and Frégnac 2003). Most of the time a decrease in the response is observed (either a decrease in the frequency of the response or/and in the number of spike emitted; see Hubel and Wiesel 1968), rather than a complete suppression as observed here. Nevertheless, phenomena of total suppression have also already been reported in physiological recordings. Goldberg and Wurtz (1972) showed a complete suppression of SCs response when increasing the size of a visual stimulus (see their Figure 4). Additionally, more recently, in a study on SCs receptive field, Wang et al. (2010) reported that the activity of SCs neurons was completely suppressed for large stimuli centred on the tested neurons (see their Figure 5). Our study brings some clues concerning mechanisms underlying these suppressive phenomena. Indeed, their neural substrates remain debated (Sachdev, Krause, and Mazer 2012). The origin of the suppression is proposed to be due to 1) a decrease of feedforward activation 2) interactions involving local lateral connections or, finally, 3) feedback connections from higher areas. The present study confirms, on a theoretical ground, that centre surround interactions in a single layer based on the most up to date physiological evidence from SC is sufficient to provide total suppression of the response for a certain range of stimulus sizes.

For any given surround suppression phenomenon, other observations in the present work provide predictions to test the hypothesis that it might be driven by short inhibitory lateral connections. First, increasing the size of a line stimulus should lead, after the suppression phase; to the reappearance of activation clusters on sites corresponding to extremities (see Figure 3). Second, this reappearance should be observed with a significant latency increase if a delayed wave of inhibition is present (see Figure 4). Third, when two stimulations are tested, maximal activity suppression should also be observed for a specific distance (see Figure 7).

4.2 Suppression phenomena: Modelling results

Similar models with MH connections have already been suggested to reproduce surround suppression (Sceniak et al. 1999; Schwabe et al. 2006; Spratling 2010). Only Sceniak et al. (1999) also showed total suppression (see their Figure 2F). Nevertheless none of them were constructed with spiking neurons. Further than the effect of the spatial organization of inhibition and excitation, our work gives an insight in how the *dynamic* of the inhibition and excitation can shape the suppression. In the present model, it is a delayed wave of inhibition – i.e. after an initial rise of membrane potential-- which drives the surround suppression. A change of the inhibition time constant would modify the suppression effect. This dynamics of the membrane potential during the surround suppression phenomenon could be investigated in experimental intracellular recordings and, if matching those observed in the present study, be used to infer the inhibition time constant of the local circuitry.

Finally, our results suggest an optimal size of visual stimuli which minimizes the latency to trigger a spiking cluster (Figure 4D) in our target selection model. This is in line with the modelling work of Marino et al. (2012) –see their figure 6F— who were working with a population rate model and an arbitrary threshold to trigger saccades. They also observed a U-shape relationship, but didn't observe a total suppression.

4.3 Suppression phenomena: Behavioral results

If the suppression phenomenon observed in our model exists in the oculomotor system, this predicts that large stimuli will lead to fewer saccades with short latency than would small stimuli. Ploner et al. (2004) observed this type of effect in a behavioural study: saccades with short latency were less numerous for large targets (10°), whereas saccades with short latency were more frequent for small target sizes (1°). More precisely concerning this latency question, the U-shape curve for the relationship between the membrane potential evolution speed and the size of the stimulus (Figure 4D, see also Figure 6F of Marino et al. 2012) is in line with the relationship shown by Boch et al. (1984) between express saccades latency and the size of the target (see their figure 5).

The observed suppression would similarly predict that larger distracting stimuli could paradoxically interfere less with saccades to a nearby target than might smaller distractors. Such a pattern was observed by Tandonnet et al. (2012). Their work focused on the Global Effect, which is the tendency for saccades to land in between to nearby visual stimuli (Findlay 1982). Using a target-distractor couple, they found a U-shaped curve for such deviation while increasing the distractor size: first the distractor is too small to have a strong influence, then its increase in size makes its influence grows, but from a given size its influence begin to decrease. This loss of weight for larger stimuli could be explained by a decreased response in a saliency map such as the SCs or the LIP. Finally, the results of Stigchel et al. (2012) consisting in a smaller extent of the global effect for large stimuli may also be explainable by a suppression of large stimuli. Note that while Tandonnet et al. (2012) observed the average shift of the landing positions, Stigchel et al. (2012) observed the split from unimodal to bimodal distribution. All these results suggest that different degrees of suppression are observable at the behavioural level. It remains to be investigated whether total suppression phenomena can also be detected – for single stimulus, and for two stimuli.

4.4 Spatial Deviation: the Fusion effect

Our DNF-MH demonstrates deviation of the spiking clusters from the initial input locations. Such deviation can be detrimental, for instance, when the DNF-MH is used as a target selection map which has to select among different points of interest sent by satellite structures. Indeed, with such deviation,

665 the selected target would not correspond to any prior points of interest. We discuss here whether these
666 deviations have already been observed at the neurophysiological, modelling or behavioural level.

667 When the model is stimulated at two nearby locations a single spiking cluster emerges in-between
668 them. The cluster is closer to the stronger stimulation location – in proportion to its relative strength
669 – and it is of the same width as spiking clusters induced by a single stimulation. This phenomenon
670 of *attraction (and fusion)* was described for the first time by Amari (1977) in a DNF-MH based on a
671 firing rate neuron model. In the context of the spatial working memory, Compte et al. (2000) proposed
672 a model consisting in a one dimensional DNF with a MH connectivity pattern. Interestingly this group
673 recently demonstrated that this model could lead to phenomena of attraction and fusion (Almeida et
674 al. 2015). The findings of the present study extend these previous observations to a 2D spiking neuron
675 networks.

676
677 On the behavioural side, the tendency for saccades to land in between two simultaneous and nearby
678 visual stimuli is known as the Global Effect or saccade averaging (Findlay 1982). Concerning the
679 neurophysiological approach, Glimcher and Sparks (1993) showed that this fusion phenomenon could
680 occur in the SCi when an intermediate saccade is made between two visual stimuli presented
681 simultaneously. Edelman and Keller (1998) added that this could be the case for saccades of latency
682 in the average range while two distinct bumps of activity would stand on the SCi for shortest latency
683 saccades. However, whether a fusion of activity in the SCs or the SCi can explain the Global Effect
684 is still matter of debate. Arai et al. (1994) implemented a saccadic system model using a DNF-MH to
685 simulate the SC layers. Their model took into account the SC spatial compression and, in their test
686 using fusion to explain the Global Effect, one can notice hypermetria (overshoot) of the output
687 saccade (see their figure 10). Katnani and Gandhi (2011) brought further insight for that result: when
688 the DNF-MH phenomenon of fusion is applied in SC space (Note that the SCs and SCi are assumed
689 to have to an equivalent mapping; cf. Schiller and Stryker 1972), this would lead systematically to
690 overshooting averaging saccades in external or retinotopic space. On the other hand, they
691 demonstrated that a vector averaging of two steady bumps of the SC space would lead neither to a
692 hypo- nor a hypermetria. They, however, note that if the phenomenon of attraction could lead to a
693 wider bump of activity (wider on the axis formed by the two input stimulations, leading to an elliptic
694 shape), the hypermetria would be corrected.

695 Recently, Vokoun et al. (2014) have reported in their work applying photodiode stimulations that on
696 a coronal slice of the *superficial layers* of the rat SC- “simultaneous stimulation of two nearby sites
697 resulted in a single, merged peak centered between the two sites”. They suggest that such a
698 phenomenon could explain the Global Effect. Importantly, they observed that an activity bump
699 induced by the simultaneous stimulation of two loci is wider than an activity bump induced by a
700 single stimulation. That results interestingly echoes to a previous behavioural study observing that
701 larger visual stimuli can lead to a wider distribution of saccade landing positions (Tandonnet and Vitu
702 2013). Under the considerations of Katnani and Gandhi (2011) this spread of activation could correct
703 the hypermetria issue discussed above, but it is important to note that a single layer bistable DNF-
704 MH model such as ours could not replicate such a spread of activity - the fused activity has the same
705 width as that for a single stimulus because the stable cluster size is set by the width of the MH.

706 4.5 Spatial Deviation: the Repulsion effect

707 When two clusters of activity were induced by two stimuli, they tended to deviate away from each
708 other (see Figure 7). Here also both the early work of Amari (1977) and the recent study of Almeida
709 et al. (2015) already observed this phenomenon in 1D models. Again our results allow to extend these
710 findings to a 2D situation. To evoke this repulsion phenomenon, Amari (1977) explains that bumps
711 of activity tend to climb up inhibition slopes. Then, the repulsion is reserved to MHs which have a
712 range short enough to allow a stimulation to “climb” the outer inhibition slope of another.

Concerning the behavioural level, Wang et al. (2012) as well as Wang and Theeuwes (2014) suggest that if this phenomenon is present in the SC, it could explain the trends of saccade trajectories to deviate away from a distractor. Wang and Theeuwes (2014) also report a shift of the landing positions away from the previous fixation stimulus when varying its timing which might be explain by repulsion. However, to the best of our knowledge, repulsion in the bimodal distribution of landing positions to two simultaneously presented stimuli or in the internal representation of stimuli position has never been observed. This may be due to the difficulty to track back a phenomenon occurring in the SCs from behavioural data. For instance, the strongest repulsion effect we observed occurred when there are two spiking clusters emerging on the map. Nevertheless, if there is vector averaging downstream, at the behavioural level only a Global effect might be observed.

Finally, at the neurophysiological level, Vokoun et al. (2014) studied activations in coronal slices of the superficial layers of the rat SC after concomitant stimulation of 2 two sites. They did not observe any repulsion (nor any suppression) effect despite the exploration of numerous distances between the two stimulated sites. Hence, even though evidences have been found recently for a local Mexican hat kernel in the SCs (Phongphanphanee et al. 2014), the lack of concordance between the present study results and Vokoun et al. (2014)'s results question if the SCs can be modelled with a simple DNF-MH (see also the end of the previous section, 4.4). However, a possible alternative to explain this lack of concordance is that the coronal slicing used by Vokoun et al. (2014) may have damaged part of the lateral inhibition system altering the MH kernel, its size and its properties. Hence, further neurophysiological works are required to shed more lights on 1) the link between fusion of activity in the SC layers and Global Effect, and 2) on what extent those natural phenomena can be modelled with a simple DNF-MH.

5 Conclusion

We constructed a DNF-MH integrating short range MH connections based on recent results obtained in the superficial layers of the SC, and we tested how it performs in very simple target selection tasks: 1) the localization of a single stimulus of different sizes; 2) the selection and localization of the strongest of a pair of stimulations.

Our work demonstrates that even a short range inhibition (i.e. only slightly larger than the excitation; ratio of 1.2) can enable a selection dynamic. However, it also highlights noticeable phenomena emerging from the model during those tasks: suppression, multi-spot selection, attraction/fusion and repulsion. If the DNF-MH is used as a target selection map as it is thought to be the case for the SCs, such attraction and repulsion would impair the spatial precision of the selection while the suppression would delay or hinder selection. In short, those properties suggest that the SCs is an imperfect winner-take-all selection system. At the same time, those properties constitute a collection of testable predictions to verify this point and the pertinence of using a DNF with short range MH to model the SCs. In parallel, future modelling work may investigate whether the phenomena we observed survive more advanced implementations of the SC dynamics. Notably, 1) when one implements the transient visual burst dynamics in SCs; 2) when one implements the SCi layer and the motor executions. Finally, results obtained in the present study have been obtained with activity in the range of what can be observed in the SC (up to 600 Hz). Further works remain to be done to explore what would be observed in DNF with lower maximum frequency.

Interestingly, attraction and repulsion phenomena have recently been reported when using DNF-MHs in spatial working memory tasks, and they have been successfully related to actual behavioural imprecisions (Almeida et al. 2015). Those results support the point that DNF-MHs are imperfect winner-take-all systems and relevant models of biological networks at the same time.

758 **Acknowledgements:**

759 This work was supported by a PhD studentship from Cardiff University, and a grant from the ESRC
760 (ES/K002325/1).

761 **References:**

762

- 763 Allman, John, Francis Miezin, and EveLynn McGuinness. 1985. "Stimulus Specific Responses from
764 beyond the Classical Receptive Field: Neurophysiological Mechanisms for Local-Global
765 Comparisons in Visual Neurons." *Annual Review of Neuroscience* 8 (1): 407–30.
- 766 Almeida, Rita, Joao Barbosa, and Albert Compte. 2015. "Neural Circuit Basis of Visuo-Spatial
767 Working Memory Precision: A Computational and Behavioral Study." *Journal of*
768 *Neurophysiology*, jn – 00362.
- 769 Amari, Shun-ichi. 1977. "Dynamics of Pattern Formation in Lateral-Inhibition Type Neural Fields." *Biological Cybernetics* 27 (2): 77–87.
- 770
- 771 Anderson, Russell W., Edward L. Keller, Neeraj J. Gandhi, and Sanjoy Das. 1998. "Two-
772 Dimensional Saccade-Related Population Activity in Superior Colliculus in Monkey." *Journal of Neurophysiology* 80 (2): 798–817.
- 773
- 774 Arai, K., E. L. Keller, and J. A. Edelman. 1993. "A Spatio-Temporal Neural Network Model of
775 Saccade Generation." In *Neural Networks, 1993., IEEE International Conference on*, 70–74.
776 http://ieeexplore.ieee.org/xpls/abs_all.jsp?arnumber=298529.
- 777 Arai, Kuniharu, Edward L. Keller, and Jay A. Edelman. 1994. "Two-Dimensional Neural Network
778 Model of the Primate Saccadic System." *Neural Networks* 7 (6–7): 1115–35.
779 doi:10.1016/S0893-6080(05)80162-5.
- 780 Behan, M., and N. M. Kime. 1996. "Intrinsic Circuitry in the Deep Layers of the Cat Superior
781 Colliculus." *Visual Neuroscience* 13 (06): 1031–42.
- 782 Boch, R., Prof Dr B. Fischer, and E. Ramsperger. 1984. "Express-Saccades of the Monkey: Reaction
783 Times versus Intensity, Size, Duration, and Eccentricity of Their Targets." *Experimental*
784 *Brain Research* 55 (2): 223–31. doi:10.1007/BF00237273.
- 785 Bompas, A., and P. Sumner. 2011. "Saccadic Inhibition Reveals the Timing of Automatic and
786 Voluntary Signals in the Human Brain." *Journal of Neuroscience* 31 (35): 12501–12.
787 doi:10.1523/JNEUROSCI.2234-11.2011.
- 788 Bretzner, Lars, and Tony Lindeberg. 1998. "Feature Tracking with Automatic Selection of Spatial
789 Scales." *Computer Vision and Image Understanding* 71 (3): 385–92.
- 790 Brunel, Nicolas, and Mark CW van Rossum. 2007. "Quantitative Investigations of Electrical Nerve
791 Excitation Treated as Polarization." *Biological Cybernetics* 97 (5): 341–49.
- 792 Compte, Albert, Nicolas Brunel, Patricia S. Goldman-Rakic, and Xiao-Jing Wang. 2000. "Synaptic
793 Mechanisms and Network Dynamics Underlying Spatial Working Memory in a Cortical
794 Network Model." *Cerebral Cortex* 10 (9): 910–23.
- 795 Edelman, Jay A., and Edward L. Keller. 1998. "Dependence on Target Configuration of Express
796 Saccade-Related Activity in the Primate Superior Colliculus." *Journal of Neurophysiology* 80
797 (3): 1407–26.
- 798 Erlhagen, Wolfram, and Estela Bicho. 2006. "The Dynamic Neural Field Approach to Cognitive
799 Robotics." *Journal of Neural Engineering* 3 (3): R36.
- 800 Findlay, John M. 1982. "Global Visual Processing for Saccadic Eye Movements." *Vision Research*
801 22 (8): 1033–45.
- 802 Glimcher, Paul W., and David L. Sparks. 1993. "Representation of Averaging Saccades in the
803 Superior Colliculus of the Monkey." *Experimental Brain Research* 95 (3): 429–35.
- 804 Goldberg, Michael E., and Robert H. Wurtz. 1972. "Activity of Superior Colliculus in Behaving
805 Monkey. I. Visual Receptive Fields of Single Neurons." *J Neurophysiol* 35 (4): 542–59.

806 Goodman, Dan, and Romain Brette. 2008. "Brian: A Simulator for Spiking Neural Networks in
807 Python." *Frontiers in Neuroinformatics* 2.
808 <http://www.ncbi.nlm.nih.gov/pmc/articles/pmc2605403/>.

809 Goodman, Dan FM, and Romain Brette. 2009. "The Brian Simulator." *Frontiers in Neuroscience* 3
810 (2): 192.

811 Guillaume, Alain, and Denis Pélisson. 2001. "Gaze Shifts Evoked by Electrical Stimulation of the
812 Superior Colliculus in the Head-Unrestrained Cat. I. Effect of the Locus and of the Parameters
813 of Stimulation." *European Journal of Neuroscience* 14 (8): 1331–44.

814 Hodgkin, A. L., and A. F. Huxley. 1952. "A Quantitative Description of Membrane Current and Its
815 Application to Conduction and Excitation in Nerve." *The Journal of Physiology* 117 (4): 500–
816 544.

817 Hubel, D. H., and T. N. Wiesel. 1968. "Receptive Fields and Functional Architecture of Monkey
818 Striate Cortex." *The Journal of Physiology* 195 (1): 215–43.

819 Indiveri, Giacomo, Elisabetta Chicca, and Rodney J. Douglas. 2009. "Artificial Cognitive Systems:
820 From VLSI Networks of Spiking Neurons to Neuromorphic Cognition." *Cognitive*
821 *Computation* 1 (2): 119–27. doi:10.1007/s12559-008-9003-6.

822 Isa, Tadashi. 2002. "Intrinsic Processing in the Mammalian Superior Colliculus." *Current Opinion in*
823 *Neurobiology* 12 (6): 668–77. doi:10.1016/S0959-4388(02)00387-2.

824 Isa, Tadashi, and William C. Hall. 2009. "Exploring the Superior Colliculus In Vitro." *Journal of*
825 *Neurophysiology* 102 (5): 2581–93. doi:10.1152/jn.00498.2009.

826 Kang, Kukjin, Michael Shelley, and Haim Sompolinsky. 2003. "Mexican Hats and Pinwheels in
827 Visual Cortex." *Proceedings of the National Academy of Sciences* 100 (5): 2848–53.
828 doi:10.1073/pnas.0138051100.

829 Katnani, Husam A., and Neeraj J. Gandhi. 2011. "Order of Operations for Decoding Superior
830 Colliculus Activity for Saccade Generation." *Journal of Neurophysiology* 106 (3): 1250–59.

831 Koch, Christof, and Shimon Ullman. 1985. "Shifts in Selective Visual Attention: Towards the
832 Underlying Neural Circuitry." In *Matters of Intelligence*, 115–41. Springer.
833 http://link.springer.com/chapter/10.1007/978-94-009-3833-5_5.

834 Kong, Hui, H.C. Akakin, and S.E. Sarma. 2013. "A Generalized Laplacian of Gaussian Filter for
835 Blob Detection and Its Applications." *IEEE Transactions on Cybernetics* 43 (6): 1719–33.
836 doi:10.1109/TSMCB.2012.2228639.

837 Kopecz, Klaus. 1995. "Saccadic Reaction Times in Gap/overlap Paradigms: A Model Based on
838 Integration of Intentional and Visual Information on Neural, Dynamic Fields." *Vision*
839 *Research* 35 (20): 2911–25. doi:10.1016/0042-6989(95)00066-9.

840 Kopecz, Klaus, and Gregor Schöner. 1995. "Saccadic Motor Planning by Integrating Visual
841 Information and Pre-Information on Neural Dynamic Fields." *Biological Cybernetics* 73 (1):
842 49–60. doi:10.1007/BF00199055.

843 Krauzlis, Richard J., Dorion Liston, and Christopher D. Carello. 2004. "Target Selection and the
844 Superior Colliculus: Goals, Choices and Hypotheses." *Vision Research* 44 (12): 1445–51.

845 Lapicque, L. 1907. "Recherches Quantitatives Sur L'excitation Electrique Des Nerfs Traitee Comme
846 Une Polarization." *J Physiol Pathol Gen* 9: 620–35.

847 Lee, P., and W. C. Hall. 2006. "An In Vitro Study of Horizontal Connections in the Intermediate
848 Layer of the Superior Colliculus." *Journal of Neuroscience* 26 (18): 4763–68.
849 doi:10.1523/JNEUROSCI.0724-06.2006.

850 Lo, Chung-Chuan, Leanne Boucher, Martin Paré, Jeffrey D. Schall, and Xiao-Jing Wang. 2009.
851 "Proactive Inhibitory Control and Attractor Dynamics in Countermanding Action: A Spiking
852 Neural Circuit Model." *The Journal of Neuroscience* 29 (28): 9059–71.
853 doi:10.1523/JNEUROSCI.6164-08.2009.

854 Lowe, David G. 1999. "Object Recognition from Local Scale-Invariant Features." In *Computer*
855 *Vision, 1999. The Proceedings of the Seventh IEEE International Conference on*, 2:1150–57.
856 Ieee. http://ieeexplore.ieee.org/xpls/abs_all.jsp?arnumber=790410.

- Marino, Robert A., Thomas P. Trappenberg, Michael Dorris, and Douglas P. Munoz. 2012. "Spatial Interactions in the Superior Colliculus Predict Saccade Behavior in a Neural Field Model." *Journal of Cognitive Neuroscience* 24 (2): 315–36.
- McIlwain, JAMES T. 1982. "Lateral Spread of Neural Excitation during Microstimulation in Intermediate Gray Layer of Cat's Superior Colliculus." *Journal of Neurophysiology* 47 (2): 167–78.
- Meeter, Martijn, Stefan Van der Stigchel, and Jan Theeuwes. 2010. "A Competitive Integration Model of Exogenous and Endogenous Eye Movements." *Biological Cybernetics* 102 (4): 271–91. doi:10.1007/s00422-010-0365-y.
- Meredith, M. Alex, and Ary S. Ramoa. 1998. "Intrinsic Circuitry of the Superior Colliculus: Pharmacophysiological Identification of Horizontally Oriented Inhibitory Interneurons." *Journal of Neurophysiology* 79 (3): 1597–1602.
- Millner, Sebastian, Andreas Grübl, Karlheinz Meier, Johannes Schemmel, and Marc-olivier Schwartz. 2010. "A VLSI Implementation of the Adaptive Exponential Integrate-and-Fire Neuron Model." In , 1642–50. http://machinelearning.wustl.edu/mlpapers/papers/NIPS2010_0425.
- Morén, Jan, Tomohiro Shibata, and Kenji Doya. 2010. "Toward a Spiking-Neuron Model of the Oculomotor System." In *From Animals to Animats 11*, 104–13. Springer. http://link.springer.com/chapter/10.1007/978-3-642-15193-4_10.
- . 2013. "The Mechanism of Saccade Motor Pattern Generation Investigated by a Large-Scale Spiking Neuron Model of the Superior Colliculus." *PLoS ONE* 8 (2): e57134. doi:10.1371/journal.pone.0057134.
- Munoz, Douglas P. 2002. "Commentary: Saccadic Eye Movements: Overview of Neural Circuitry." *Progress in Brain Research* 140: 89–96.
- Munoz, Douglas P., and Peter J. Istvan. 1998. "Lateral Inhibitory Interactions in the Intermediate Layers of the Monkey Superior Colliculus." *Journal of Neurophysiology* 79 (3): 1193–1209.
- Phongphanphanee, Penphimon, Robert A. Marino, Katsuyuki Kaneda, Yuchio Yanagawa, Douglas P. Munoz, and Tadashi Isa. 2014. "Distinct Local Circuit Properties of the Superficial and Intermediate Layers of the Rodent Superior Colliculus." *European Journal of Neuroscience*, April, n/a – n/a. doi:10.1111/ejn.12579.
- Ploner, Christoph J., Florian Ostendorf, and Sandra Dick. 2004. "Target Size Modulates Saccadic Eye Movements in Humans." *Behavioral Neuroscience* 118 (1): 237–42. doi:10.1037/0735-7044.118.1.237.
- Richter, M., Y. Sandamirskaya, and G. Schoner. 2012. "A Robotic Architecture for Action Selection and Behavioral Organization Inspired by Human Cognition." In *2012 IEEE/RSJ International Conference on Intelligent Robots and Systems (IROS)*, 2457–64. doi:10.1109/IROS.2012.6386153.
- Robinson, D. A. 1972. "Eye Movements Evoked by Collicular Stimulation in the Alert Monkey." *Vision Research* 12 (11): 1795–1808.
- Sachdev, Robert NS, Matthew R. Krause, and James A. Mazer. 2012. "Surround Suppression and Sparse Coding in Visual and Barrel Cortices." *Frontiers in Neural Circuits* 6. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3389675/>.
- Sandamirskaya, Yulia. 2014. "Dynamic Neural Fields as a Step toward Cognitive Neuromorphic Architectures." *Frontiers in Neuroscience* 7. doi:10.3389/fnins.2013.00276.
- Sceniak, Michael P., Dario L. Ringach, Michael J. Hawken, and Robert Shapley. 1999. "Contrast's Effect on Spatial Summation by Macaque V1 Neurons." *Nature Neuroscience* 2 (8): 733–39.
- Schiller, Peter H., and Michael Stryker. 1972. "Single-Unit Recording and Stimulation in Superior Colliculus of the Alert Rhesus Monkey." *Journal of Neurophysiology* 35 (6): 915–24.
- Schwabe, Lars, Klaus Obermayer, Alessandra Angelucci, and Paul C. Bressloff. 2006. "The Role of Feedback in Shaping the Extra-Classical Receptive Field of Cortical Neurons: A Recurrent Network Model." *The Journal of Neuroscience* 26 (36): 9117–29. doi:10.1523/JNEUROSCI.1253-06.2006.

- Series, Peggy, Jean Lorenceau, and Yves Frégnac. 2003. "The 'silent' Surround of V1 Receptive Fields: Theory and Experiments." *Journal of Physiology-Paris* 97 (4): 453–74.
- Shadlen, Michael N., and William T. Newsome. 1998. "The Variable Discharge of Cortical Neurons: Implications for Connectivity, Computation, and Information Coding." *The Journal of Neuroscience* 18 (10): 3870–96.
- Sparks, David L. 1986. "Translation of Sensory Signals into Commands for Control of Saccadic Eye Movements: Role of Primate Superior Colliculus." *Physiological Reviews* 66 (1): 118–71.
- . 2002. "The Brainstem Control of Saccadic Eye Movements." *Nature Reviews Neuroscience* 3 (12): 952–64.
- Sparks, David L., Richard Holland, and Barton L. Guthrie. 1976. "Size and Distribution of Movement Fields in the Monkey Superior Colliculus." *Brain Research* 113 (1): 21–34. doi:10.1016/0006-8993(76)90003-2.
- Spratling, Michael W. 2010. "Predictive Coding as a Model of Response Properties in Cortical Area V1." *The Journal of Neuroscience* 30 (9): 3531–43. doi:10.1523/JNEUROSCI.4911-09.2010.
- Tandonnet, Christophe, Delphine Massendari, and Françoise Vitu. 2012. "When Larger Visual Distractors Become Less Disruptive: Behavioral Evidence for Lateral Inhibition in Saccade Generation." *Journal of Vision* 12 (4): 2. doi:10.1167/12.4.2.
- Tandonnet, Christophe, and Françoise Vitu. 2013. "Stimulus Properties and Saccade Metrics: When Local Features Are More Critical than Global Features." *Behavioral Neuroscience* 127 (1): 121.
- Trappenberg, Thomas P., Michael C. Dorris, Douglas P. Munoz, and Raymond M. Klein. 2001. "A Model of Saccade Initiation Based on the Competitive Integration of Exogenous and Endogenous Signals in the Superior Colliculus." *Journal of Cognitive Neuroscience* 13 (2): 256–71.
- Van der Stigchel, S., J. Heeman, and T.C.W. Nijboer. 2012. "Averaging Is Not Everything: The Saccade Global Effect Weakens with Increasing Stimulus Size." *Vision Research* 62 (June): 108–15. doi:10.1016/j.visres.2012.04.003.
- van der Walt, Stéfan, S. Chris Colbert, and Gaël Varoquaux. 2011. "The NumPy Array: A Structure for Efficient Numerical Computation." *Computing in Science & Engineering* 13 (2): 22–30. doi:10.1109/MCSE.2011.37.
- Vokoun, Corinne R., Xin Huang, Meyer B. Jackson, and Michele A. Basso. 2014. "Response Normalization in the Superficial Layers of the Superior Colliculus as a Possible Mechanism for Saccadic Averaging." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 34 (23): 7976–87. doi:10.1523/JNEUROSCI.3022-13.2014.
- Wang, Lupeng, Rashmi Sarnaik, Krsna Rangarajan, Xiaorong Liu, and Jianhua Cang. 2010. "Visual Receptive Field Properties of Neurons in the Superficial Superior Colliculus of the Mouse." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 30 (49): 16573–84. doi:10.1523/JNEUROSCI.3305-10.2010.
- Wang, Zhiguo, Wouter Kruijne, and Jan Theeuwes. 2012. "Lateral Interactions in the Superior Colliculus Produce Saccade Deviation in a Neural Field Model." *Vision Research* 62 (June): 66–74. doi:10.1016/j.visres.2012.03.024.
- Wang, Zhiguo, and Jan Theeuwes. 2014. "Distractor Evoked Deviations of Saccade Trajectory Are Modulated by Fixation Activity in the Superior Colliculus: Computational and Behavioral Evidence." *PLoS ONE* 9 (12): e116382. doi:10.1371/journal.pone.0116382.
- Wilimzig, Claudia, Stefan Schneider, and Gregor Schöner. 2006. "The Time Course of Saccadic Decision Making: Dynamic Field Theory." *Neural Networks, Neurobiology of Decision Making Neurobiology of Decision Making*, 19 (8): 1059–74. doi:10.1016/j.neunet.2006.03.003.
- Zeng, Chi, Yongjie Li, and Chaoyi Li. 2011. "Center-surround Interaction with Adaptive Inhibition: A Computational Model for Contour Detection." *NeuroImage* 55 (1): 49–66. doi:10.1016/j.neuroimage.2010.11.067.

Provisional

Figure 1.TIF




Figure 2.TIF




Figure 3.TIF




Figure 4.TIF




Figure 5.TIF




Figure 6.TIF




Figure 7.TIF




Figure 8.TIF

